### XMM-Newton CCF Release Note

#### XMM-CCF-REL-386

## OM Grisms Calibration: time dependent sensitivity correction

S. Rosen

December 3, 2021

## 1 CCF components

| Name of CCF      | VALDATE             | EVALDATE | List of Blocks changed | XSCS flag |
|------------------|---------------------|----------|------------------------|-----------|
| OM_GRISMCAL_0007 | 2000-01-01T00:00:00 | _        | TDS_CORR               | NO        |

# 2 Changes

#### 2.1 Time dependent sensitivity degradation correction

The sensitivity of the Optical Monitor (OM) on board XMM-Newton is affected by a time dependent degradation. This is due to three main effects: the degradation of the S-20 photocathode, contamination, and the aging of the MCP. The first two of these effects are known to be wavelength dependent

This time dependent degradation is described, for the colour filters (V, B, U, UVW1, UVM2, UVW2) in XMM-SOC-CAL-TN-0207, where a correction factor is defined and computed. A corresponding description of the analysis of time-dependent degradation in OM grism data was provided in XMM-SOC-CAL-TN-0218 and is updated in XMM-SOC-CAL-TN-0222. Note that the tabulated degradation functions and correction tables shown in XMM-SOC-CAL-TN-0222, which pertain to OM\_GRISMCAL\_0006.CCF, are superseded by the values in this release note.

#### 2.2 The correction

In OM\_GRISMCAL\_0005.CCF, an extension containing the corrections for the time dependent sensitivity degradation of both OM grisms, was introduced for the first time, in conjunction with release

UV Grism Year V Grism2000 1.000 1.000 2002 1.011 1.009 2004 1.023 1.018 2006 1.035 1.027 2008 1.047 1.036 2010 1.060 1.046 2012 1.073 1.055 2014 1.086 1.065 2016 1.100 1.075 2018 1.114 1.085 2020 1.128 1.096 2022 1.1421.106

Table 1: Correction factors for the time dependent degradation of the OM grisms.

note XMM-CCF-REL-359. A subsequent update (OM\_GRISMCAL\_0006.CCF, XMM-CCF-REL-377) was released in 2020.

1.157

1.117

2024

Here we provide a new update the OM grism calibration CCF, based on the latest available OM grism calibration data derived from the routinely observed spectrophotometric standard stars, BPM16274, GD 153 and Hz2. The latest data included in the analysis come from XMM-Newton revolution 3955 (July 2021).

The correction factors for each grism, derived from the inverse of the functions fitted to the data shown in figure 1, are given in Table 1 on a uniformly spaced grid of observation epochs, with extrapolation to 2024. Since SAS 18.0, the fluxed OM grism spectra of sources processed with the XMM-Newton SAS, include a correction, interpolated from the correction factors in the CCF used at the time of processing. Note, also, that the count rates in the same spectral files are not scaled. The correction factor applied to a given fluxed spectrum is stored in the FITS keyword **TDS\_CORR**.

# 3 Scientific Impact of this Update

These corrections allow users to compare the fluxes of any source observed with the OM grisms at different epochs. These corrections extend the range of application to 2024.0.






Figure 1: Normalised measured count rates in various spectral bands from the spectra of BPM16274 (blue), GD153 (green) and Hz2 (red) for the UV grism (upper panel) and Visible grism (lower panel). The solid black lines are the linear fits to the decline in the current analysis while the dashed black lines are the fits from the previous analysis that went into OM\_GRISMCAL\_0006. For each grism, the results are obtained via a simultaneous linear fit to all the wavelength band measurements from all 3 stars, where the slope is common to all stars and wavelength bands but the normalisations, computed at 2000.0, are star and band dependent (see XMM-SOC-CAL-TN-0222).

### 4 Estimated Scientific Quality

Absolute fluxes obtained with the OM grisms will be corrected for degradation. The data exhibit significant scatter but 1- $\sigma$  uncertainties on the corrections of 2.3% (UV) and 1.9% (VIS) are estimated at 2024.0. These uncertainties do not include any systematics. For each grism, the fits are determined from the ensemble of valid data from all three standard stars and in all 6 wavelength bands - no resolution by wavelength band is currently available.

## 5 Expected Updates

The table of correction factors covers the period since launch until the year 2024.0. An update will be necessary for observations obtained after this date.

## 6 Test procedures

The new CCF was validated for content and format validity. The correction has been tested on grism data of the OM standard star, BPM16274, obtained in XMM-Newton revolution 3462, re-processed using **omgchain** v1.12, which employs **omgrism** v1.29, from SAS 19.0.

# 7 Summary of the test results

The fluxed spectra, based on use of the OM\_GRISMCAL\_0006 and OM\_GRISMCAL\_0007, were compared and found to be in accord with the changed correction factors. The TDS\_CORR header keyword values were verified as correct.

### References